

# Kaiser Permanente National Implant Registries



2017 Annual Report

# Table of Contents

Leadership message	3
What we provide	5
<b>Accomplishments and collaborations</b>	
Registries by the Numbers	9
ACLR Registry Leads to Reduction in Allograft Use	10
Cardiac Device Facilitating New Technology Decision Making	11
Post Market Surveillance of Stent Graft Implants	12
Using the Hip Fracture Registry for Benchmarking Patient Care	13
Monitoring Length of Stay for Shoulder Procedures	14
Opioid Use Before and After Total Joint Replacement	15
<b>Highlights from our registries</b>	
Anterior Cruciate Ligament Reconstruction	17
Cardiac Device	19
Endovascular Stent Graft	21
Hip Fracture	22
Shoulder Arthroplasty	24
Spine	26
Total Joint Replacement	28
Staff	31
Publications	32

# Leadership message

Dear Colleagues,

It is an honor and a privilege to present the 2017 **Kaiser Permanente National Implant Registries** annual report highlighting significant achievements in surgical outcomes, clinical research findings, and quality improvement initiatives. Our registries monitor patient characteristics, surgical approaches, implant characteristics, and clinical outcomes for more than 2.2 million cardiac, neurosurgery, orthopedic, and vascular implants for the lifetime of our 12 million members.

The registries' longitudinal tracking of implant performance provides the foundation for optimal surgical care through identification and dissemination of clinical best practices, translating registry findings into clinical care, and continuous monitoring of patient outcomes. We are proud to announce that our impact on quality of care was recognized in 2018 with the prestigious **Orthopaedic Research and Education Foundation Clinical Research** award for outstanding Anterior Cruciate Ligament reconstruction (ACLR) research. Our ACLR registry's translation of research findings into clinical care are highlighted in this year's annual report along with other key implant registry accomplishments:

- Decreased use of high-risk allografts and decreased use of allografts in young ACLR patients.
- Identification of medical devices at higher risk for complications and revisions
- Reduction of elapsed time from the time a hip fracture patient is admitted to the hospital to the time they have surgery.
- Increased use of total shoulder arthroplasty same-day discharge as a way to improve outcomes, enhance the patient experience, and reduce health care costs.
- Implementation of strategies to reduce unsafe prolonged opioid use among total joint replacement patients, with a marked reduction of opioid use for more than 90 days after surgery.

The success of our implant registries program is a reflection of the excellent Kaiser Permanente physician leadership and strong evidence-based medicine culture transforming quality of care. Our implant registries are enhancing quality and patient safety not only for our members but nationally and worldwide as we help set the standards for clinical best practices and implant performance.

**Liz Paxton, MA**

*Director, National Implant Registries*

**Tadashi Funahashi, MD**

*Chair, Inter-Regional Implant Registries Committee*

# Inter-Regional Implant Registries Committee Members

## **IIRC Membership**

**Scott H. Adelman, MD, FACC**

*Chair-Cardiology Core Group/National Product Council  
Chair-NCal Cardiovascular Technology Committee  
Senior Technology Advisor-Innovation and Advanced Technology*

**Ralph G. Brindis, MD, MPH, MACC, FSCAI**

*Clinical Professor of Medicine  
Department of Medicine & the Philip R. Lee Institute for Health Policy Studies  
University of California, San Francisco*

**Eric Cain, MD, MBA**

*Physician-in-Chief, TPMG  
Greater Southern Alameda Area*

**Patrick Courneya, MD**

*Executive Vice President, Hospitals  
SVP, Hospitals, Quality and Care Delivery Excellence  
Chief Medical Officer*

**Christopher M. Grant**

*Chief Operating Officer  
Permanente Federation, Program Offices*

**Jo Carol Hiatt, MD, MBA**

*Chair, National Product Council  
Assistant Medical Director, SCPMG Business Administration*

**Bradley B. Hill, MD, FACS**

*Chair, TPMG Vascular Surgery Chiefs Group  
Chief of Vascular Surgery*

**Margaret Mentakis, MD, FACS**

*Department of Surgery, South Sacramento  
KPHC Procedural and Perioperative Services  
TPMG Technology Leader*

**Ronald Navarro, MD**

*Asst. Area Medical Director Surgical Services  
Inter-regional Orthopedic Chief*

**Murray N. Ross, Ph.D.**

*Vice President and Director, Institute for Health Policy  
Kaiser Foundation Health Plan Enhancing Patient Safety and Quality of Care*

# What we provide

Registries play a critical role in enhancing quality of care by identifying variation and clinical best practices and providing feedback to frontline staff and clinicians using a variety of dynamic feedback mechanisms.





# Accomplishments and collaborations

## Kaiser Permanente Registry Work Receives National Recognition

Permanent Medicine physicians and Kaiser Permanente ACL Reconstruction Registry (KPACLR) have been recognized by the American Academy of Orthopaedic Surgeons with the prestigious Orthopaedic Research and Education Foundation (OREF) Clinical Research Award.

Translating registry findings into evidence-based clinical practice is the goal of a clinical registry. Furthermore, a registry's success heavily relies on continued participation from physicians, as well as the administrative team who organize, validate, and communicate the findings. Kaiser Permanente currently maintains eight such registries: cardiac, vascular, and six in orthopedics (total hip, total knee, ACLR, shoulder, hip fracture, spine). By tracking patients and procedures, measuring which factors improve or are detrimental to patient care, and then sharing this information with physicians, the Registry program has identified a direct correlation to improved outcomes over time.

Gregory Maletis, MD, Lead Physician for the Kaiser Permanente ACLR Registry, emphasizes the impact of registries on patient care: "With most research it is difficult to determine whether the findings from a specific study actually change the way clinicians practice. Registries are unique in that one can identify risk factors for good or poor outcomes and provide that information to clinicians. It is then possible to track the positive change in practice patterns which lead to improved patient outcomes."

The Registry program at Kaiser Permanente was inspired by Sweden's own healthcare registries that trace health data spanning many decades. Given Kaiser Permanente's integrated care model and ongoing encouragement of physician collaboration, physicians leaders believed that KP was well positioned to model these similar efforts and build a registry internally.




Gregory Maletis, MD was well positioned to model these similar efforts and build a registry internally.

The Kaiser Permanente ACL Reconstruction Registry (KPACLR) is one such example at Kaiser Permanente that has allowed Permanent Medicine physicians to continually improve the treatment of patients with anterior cruciate ligament (ACL) injuries. With over a decade of data collection and analysis, KP clinicians and patients have contributed to making the KPACLR the largest ACL registry in the United States and have successfully demonstrated its direct impact on the quality of outcomes for patients.

The ACLR Registry work was recently acknowledged by the American Academy of Orthopaedic Surgeons in the form of the prestigious Orthopaedic Research and Education Foundation (OREF) Clinical Research Award.



Tadashi Funahashi, MD, Assistant Regional Medical Director and Chair, Kaiser Permanente Inter-Regional Implant Registries, reflected on the award and the journey to get there: "We founded the Registries with the aspiration of building a reputation for Kaiser Permanente Orthopedic Surgery that brought pride to every orthopedic surgeon. The foundational idea was that collaboration and integration, a hallmark of Permanent Medicine, could create a world class orthopedic registry. After over a decade of effort and contributions, this recognition is shared by every orthopedic surgeon, and is a tremendously proud moment for all of us."



Every patient across eight KP regions who undergoes an ACLR is entered in the Registry, and is prospectively monitored using strict privacy controls. In addition to standardized Registry documentation, the ACLR Registry uses EHR, claims data, and other existing administrative databases, to capture patient demographics, comorbidities, anthropometric measures, surgical procedure detail, implant information, and outcomes.

This information is validated and analyzed by the Registry team, who then pass the findings through a variety of feedback mechanisms designed to influence clinical practices of orthopedic surgeons. To aid in behavior change, surgeons receive confidential individualized reports with benchmarking metrics they can use to compare their personal activity to that of the medical center, region, and organization. In addition, quality reports, annual registry reports, comparative effectiveness studies, risk calculators, and many other resources are also made available.

In disseminating feedback to physicians, the KPACLR team has, in multiple instances, demonstrated behavior change that has led to improved quality of care and outcomes. For example, analysis of Registry data on graft choice for ACL Reconstruction demonstrated that allograft use was associated with a three times higher risk of revision compared to the use of bone-patellar tendon-bone (BPTB) autografts. Furthermore, the analysis indicated that patients  $\leq 21$  years of age were at an elevated risk of revision



if allograft tissue was employed. This information was spread to surgeons and ultimately led to a decrease in both general allograft use (45%, 2010; 33%, 2015) and in allograft use for high-risk patients (28%, 2009; 9%, 2015).

The benefits of this work are not limited to KP patients. The potential to improve outcomes for patients receiving care elsewhere exists in the tools developed and published externally. For example, the KPACLR team has published a risk calculator accessible to anyone that helps predict which graft will have the best survival rate, and therefore allows for any ACLR patient to benefit from over a decade of longitudinal data cultivated by KP's Registry team.

When discussing what other organizations can learn from KP's Registry experience, Liz Paxton, Director of the Kaiser Permanente National Implant Registries, highlights physician engagement as a critical component to the success of any registry: "Physician involvement, conceptualization, and ownership have all been key to getting our physicians to trust and act on the data. With the support of our electronic health record, we have been able to integrate additional documentation needs into the current work flow and thus reduce additional burdens on staff. Even though every contribution is relatively small or brief in nature, the cumulative impact has allowed us to create the infrastructure that has resulted in meaningful research and insights."

Source: Carly Marker, Communications Consultant, Quality



### Collaborations

- Full member of International Society of Arthroplasty Registries (ISAR). <http://www.isarhome.org/>
- IIRC contribution to the planning board and Medical Device Registries Task Force resulting in key recommendations for the FDA for national medical device surveillance system.
- Orthopedic Coordinated Registries Network (Ortho CRN): collaboration with other US registries to create a mechanism for national post-market surveillance.
- Led International Collaboration of Orthopedic Registries (ICOR) along with Cornell to develop a network of international registries for medical device surveillance. <http://www.icor-initiative.org/>
- Medical Device Epidemiology Network (MDEpinet): Ongoing collaboration with FDA and regulators, academia, other health systems, and industry to build a national medical device evaluation system by enhancing the current infrastructure for real-world data, development of innovative methodologies, and conducting clinical studies ([mdepinet.org](http://mdepinet.org)).
- Build on international collaborations with Anterior Cruciate Ligament Reconstruction (ACLR) and Shoulder Arthroplasty Registries to detect surgical best practices based upon implant selection and surgical techniques to help deliver the highest quality of care.



## National Implant Registries: By the numbers

8 regions, 9 states representing  
11.7 million members

50 hospitals

133 publications in peer-reviewed journals  
15 in 2016

170 posters & presentations at national &  
international symposia

2,400 physicians participating in implant registries

2,584 patients with enhanced surveillance due to 11  
recalls in 2015-2016

445,000+ patients in our registries

550,000+ procedures tracked for the patient's lifetime

2.15 million implants registered



## ACLR Registry feedback leads to reduction in allograft use



A sprain or tear of the anterior cruciate ligament (ACL) is one of the most common knee injuries especially among athletes who participate in sports such as soccer, football, and basketball. Some people can cope with an ACL deficient knee, but those wishing to return to pivoting and twisting sports often require ACL reconstruction (ACLR).

Many factors may contribute to the outcome of ACLR surgery, including graft choice. The two main graft options available to patients for ACL reconstruction consist of autografts (donated from the patient themselves) and allografts (donated from human cadaver tissue).

The Kaiser Permanente National ACLR Registry tracks the outcome of all allograft and autograft cases performed by KP surgeons across the United States.

By 2008, the registry showed that the annual proportion of ACLR cases using an allograft was on the rise at Kaiser Permanente, hitting a peak of 45% by 2010.

That upward trend began to change in 2012 when registry findings showed a three times higher risk of revision if an allograft was used rather than a bone-patellar tendon-bone (BPTB) autograft. In subsequent studies of allografts, the registry identified poorer results with BPTB allografts compared to soft tissue allografts. Data also showed that patients  $\leq 21$  years


of age were at increased risk for revision when allograft tissue was used.

Findings were widely disseminated to surgeons using a variety of means including peer-reviewed publications; internal and external meetings and conferences; newsletters of study findings; risk calculators; and confidential individualized reports of surgeons' outcomes. In addition, surgeon champions set a quality improvement goal to reduce allograft usage overall and specifically to decrease the use of high-risk grafts and usage in high-risk patient groups. "By getting this information out to surgeons, we have seen decreased use of high-risk allografts and decreased use of allografts in young patients," said Greg Maletis, MD, chief of orthopedics, Baldwin Park Medical Center and ACLR physician lead. "These changes have led to better outcomes for our patients with a decreased risk of revision."

Registry data also showed that BPTB autografts have the lowest revision risk. "As a result, I am doing more BPTB autografts than in the past," said Maletis. "The registry has definitely changed my practice and that of my colleagues as well."

Kaiser Permanente has been awarded the 2018 Orthopaedic Research and Education Foundation Clinical Research Award for its ACLR Registry work.

# Cardiac Device Facilitating New Technology Decision Making



Implant registries provide product performance and utilization data to the National Product Council, Kaiser Permanente’s device procurement arm, to support the selection of the highest quality implant products and technologies. Innovation and new technology are encouraged, but only to the extent that they produce improved patient outcomes.



“Before any decisions are made regarding new products, we can use the registry to see if the proposed benefits would be realized in our own KP population, rather than relying on third party information or even research that may not be valid either for our patient population or our health care delivery model,” said Nigel Gupta, MD, director, Regional Cardiac Electrophysiology Services, Los Angeles Medical Center.

Dr. Gupta serves as chair of the Cardiac Rhythm Management, Standards and Sourcing Team (CRM-SST), one of many SSTs which advise the National Product Council on contracting decisions. “The primary mission of the SSTs is to govern selection decisions on products and services, always keeping the patient in mind as our first goal, with value as an

added benefit,” said Dr. Gupta. “The registry can, at times, reveal information to us that is the deciding factor for a contract or vendor.”

The Cardiac Rhythm Management group used registry data in considering procurement of a “leadless” pacemaker that promised a reduction in infection rates when compared with traditional pacemakers.

“The fact that the leadless pacemaker studies quoted a dramatic reduction in infection rates between “historical” infection rates and those using the new pacemaker made it tempting,” said Dr. Gupta. “However, when we pulled our own data from the registry we realized that our actual rates of infection were already extremely low, approaching those achieved by the new technology, even without using the new technology.”

Registry data showing little difference in the infection rates between leadless and traditional pacemakers was instrumental in the decision to hold off adopting the new technology. “This persuaded us as a group to delay further large deployment until we could realize other benefits to merit the cost and risks of the leadless device,” said Dr. Gupta. “Without the registry data we would not have realized the potential benefit of our current position.”

## Post Market Surveillance of Stent Graft Implants



In the rare event of a medical device recall or safety advisory, the registries enable Kaiser Permanente to quickly identify all patients with that device and bring them in for appropriate management.

The Kaiser Permanente Product Recall Department contacts the registries when informed of a recall or advisory. “Each surgeon is then contacted regarding his or her own patients, so the appropriate action can be taken,” said orthopedic surgeon Mark Dillon, MD, Sacramento Medical Center. “In this way, our registry serves as an additional safety net for our patients and, as a surgeon, it makes me feel more confident in the choices I make in the operating room.”

Since the implant registries were first initiated in 2001, they have been instrumental in helping to manage 78 implant recalls and advisories, affecting 106,263 Kaiser Permanente members nationwide.

Among the safety advisories received by Kaiser Permanente in 2016 was one issued regarding a Type III endoleak in an endovascular AAA (abdominal aortic aneurysm) system. An endoleak is a complication that occurs when blood flows into the aneurysm cavity from an aorta treated with a stent, thereby increasing

the risk of aneurysm rupture and death. At the time of the advisory, several hundred systems had been implanted in patients across all KP regions.

“The registry proved invaluable in identifying patients at risk,” said Bradley Hill, MD, Chief of Vascular



Surgery, Santa Clara Medical Center. “We used the list to reach out to all the patients and make sure they were getting the surveillance they needed and had a management plan in place.” Patients with the system now receive routine, clinically appropriate imaging based on physician expertise to monitor device integrity and detect an endoleak, should it develop.

“This device problem has enabled us to fully utilize the registry for its intended purpose – to ensure the safety of our patients with AAA endografts,” said Dr. Hill. “Our patients are followed longitudinally by the registry as long as they remain KP members. As a result, our registry follow-up duration and data quality tend to be better than most other registries.”

## Using the Hip Fracture Registry for Benchmarking Patient Care



**K**aaiser Permanente registries improve quality of care by providing feedback to physicians. Registry input allows individual surgeons and medical centers to benchmark or compare their performance against others within Kaiser Permanente and, if needed, adjust their practices to improve. “The registries are very useful for clinicians, and they’re also very useful for patients and their families,” said orthopedic surgeon, Gary Zohman, MD, Anaheim Medical Center. “People want to know about infection rates or complications or the time it takes to get someone into surgery after a hip fracture — and that’s something registry data can tell them.”



The elapsed time from the time a patient is admitted to the hospital for a fractured hip to the time they have surgery is among the data captured by Kaiser Permanente’s hip fracture registry. Hip fractures can lead to a host of other serious and life-threatening conditions in the elderly population where they most often occur. “Research has shown that the best way to reduce complication and mortality rates is to stabilize a hip fracture as soon as possible after the patient has been medically optimized,” said Dr. Zohman. “That’s why we made the time from admission to surgery as one of our benchmarks.”

Kaiser Permanente established a benchmark of 48 hours as the target time for getting hip fracture

patients into surgery. “As it turns out, registry data showed us that the vast majority of our medical centers were getting 91% of our hip fracture patients into surgery within 48 hours,” noted Dr. Zohman. “The question now was, can we do even better.”

The hip fracture registry, like all registries, enables each medical center to clearly see their performance in relation to the 48-hour benchmark, the first step in identifying opportunities for improvement. “Right now, we’re at the point of showing our clinicians whether they are meeting the benchmark and help them determine if not, why not,” said Dr. Zohman. “We don’t have those results yet because this is such a new benchmark but we do have everyone focused on the goal and helping us to smooth the process.”

Getting people into emergency surgery for hip fractures is a complex process that involves the emergency department, the medical team optimizing the patient for surgery, and the operating room itself. The registry will help identify practices at each step of the way that either delay or speed the process from admission to surgery.

“It can be very unsettling when there are delays in stabilizing a hip fracture and families are very appreciative when things are done promptly,” said Dr. Zohman. “We want families to know that Kaiser Permanente is looking after their loved ones and doing everything we can to fix those hips as quickly as it’s safe to do so.”

## Monitoring Length of Stay for Shoulder Procedures



**A**lthough shoulder joint replacement is currently less common than knee or hip replacement, its use is increasing, especially as more elderly patients are becoming candidates for the procedure. In the U.S., the number of total and partial shoulder replacements increased from about 18,000 in 2000 to more than 45,000 in 2013, according to the American Academy of Orthopaedic Surgeons.

With the rate of total shoulder arthroplasty (TSA) rising, same-day discharge is gaining interest as a way to improve outcomes, enhance the patient experience, and reduce health care costs.



Kaiser Permanente shoulder surgeons have been progressively shortening the length of stay (LOS) following shoulder replacement surgery. The LOS was around 2 days from 2010 to 2013; it declined to 1+ days in the 2013 to 2016 timeframe. "We have now begun sending some

patients home the same day and wanted to assess the practice with the rigor that a peer-reviewed study would provide," said Ronald Navarro, MD, Regional Coordinating Chief of Orthopedic Surgery, South Bay Medical Center.

The Kaiser Permanente Shoulder Arthroplasty Registry conducted a study to compare same-day discharge patients with those staying one or more nights. The study showed that same-day discharge following shoulder replacement is not inferior to longer length of stay with regards to 30-day emergency visits and 90-day readmission.

Study results have contributed to an increase in same-day discharge for shoulder arthroplasty. "Same-day discharge is not yet standard practice nor do we anticipate same-day discharge for all shoulder patients," said Dr. Navarro. "We are currently testing a same-day discharge program that we will soon be rolling out to a wider audience."

Kaiser Permanente supports home recovery for same-day patients using a national program called Enhanced Recovery after Surgery (ERAS) that includes multimodal pain medications, gastric attention to avoid nausea, and attention to IV fluid volume administration to avoid bladder fullness.

"We have shown that same-day total shoulder arthroplasty is a safe and economical procedure," said Dr. Navarro. "This information provides evidence that same-day shoulder arthroplasty can provide equivalent safety and financial savings for any health care system."

## Opioid use before and after total joint replacement




Total knee and hip replacement surgeries are among the most common and effective treatments for patients with impaired function and severe pain in these joints. Opioids have traditionally been, and remain, a component in postoperative pain relief for total joint replacement (TJR) patients as a part of perioperative multimodal pain protocols.

But while opioids can be a valid course of treatment to manage acute pain, they carry the risk of addiction and overdose. Starting in 2010, Kaiser Permanente implemented a Safe and Appropriate Opioid Prescribing Program to reduce over-prescribing of opioids.

As part of ongoing clinical care improvement efforts and to support safe and appropriate prescribing of opioids for TJR patients, Kaiser Permanente's TJR Registry conducted some of the first studies looking at the effects of preoperative and postoperative opioid use on TJR patients. These studies identified how many patients were using opioids and factors associated with an increased risk for prolonged opioid use.

The studies showed that preoperative opioid use is associated with prolonged opioid use after TJR. The studies also found that continued opioid use after the first 90-day postoperative period is associated with a higher revision risk and warrants close patient follow-up.

These findings have led to the implementation of strategies at Kaiser Permanente to reduce unsafe opioid use among TJR candidates and recipients.



"We have identified the cessation of opioids 90 days after TJR as one of the process improvement goals at our medical center," said orthopedic surgeon Robert Namba, MD, Irvine Medical Center. "There is now increased awareness among our surgeons that patients requesting refills over 90 days after surgery merit closer follow-up."


At the Irvine Medical Center, dissemination of study results has led to a marked reduction of prolonged opioid use. Between 2015 and May of 2017 at the medical center, total hip replacement patients using opioids more than 90 days after surgery declined from 34% to 12%. During the same time period, total knee replacement patients using opioids more than 90 days after surgery dropped from 39% to 19%.

"All healthcare providers are challenged with addressing the opioid epidemic in the U.S.," said Dr. Namba. "Excess dispensing of opioids is not only a risk factor for our TJR patients, but also increases the quantity of opioids which may be taken and disseminated unwittingly to the friends and families of our patients."





# ACLR Registry Overview



## Description:

The anterior cruciate ligament reconstruction (ACLR) registry was established in 2005 and tracks implants and outcomes of ACLR cases. As of year-end 2016, there were 40,666 cases in the ACLR registry.

## Clinical Findings

- In our cohort of 10,190 primary ACLRs using allografts, 8425 (82.7%) received a processed allograft and 1765 (17.3%) received a non-processed allograft. The overall incidence of deep infection after ACLR was very low (0.15%). No difference in the likelihood of infection between processed and non-processed allografts could be identified. Allograft tissue is often irradiated and/or chemically processed. Studies have identified an association between this processing and a higher risk for revision; whether processing decreases infection risk has yet to be determined.
- In our cohort of 4,087 primary ACLRs, with no meniscal injury at the time of index surgery and contralateral knee normal, there was a 3.73 times higher likelihood of subsequent meniscal surgery

in the index knee compared with the contralateral knee. Compared to BPTB autografts, allografts and hamstring autografts were risk factors for subsequent meniscal surgery in the index knee. This information should be considered when determining the appropriate graft choice for ACL reconstruction, and meniscal surgery after ACL reconstruction should be considered as an outcome variable when comparing various ACL reconstruction techniques.

- When BPTB allograft tissue was used for ACLR, an overall 4.54 times adjusted higher risk of revision was observed compared with surgery performed with a BPTB autograft. Whether the tissue was irradiated with either high- or low-dose radiation, chemically processed, or not processed at all made little difference in the risk of revision. The differences in the revision risk were also consistent in younger and older patients. Surgeons and patients should be aware of the increased risk of revision when a BPTB allograft is used for ACLR.

**Registry Champions:** Gregory Maletis, MD; Tadashi Funahashi, MD; Anita Rao, MD; Mark Shaieb, MD; Ron Wyatt, MD; Terrill Julien, MD; Anne Denys, MD; Mark Davies, MD; Darin Allred, MD

# Anterior Cruciate Ligament Reconstruction Registry

## Anterior Cruciate Ligament Reconstruction KP compared to benchmarks

Characteristic n (%)	Kaiser Permanente	Danish Cruciate Ligament Register	Norwegian National Knee Ligament Register	Swedish National ACL Register	UK National Ligament Register
Start Date	Feb-05	Jul-05	Jun-04	Mar-05	Dec-12
Total N	40113	30714	25624	40252	9002
Primaries	35524 (88.6)	26457 (86.1)	23337 (91.1)	37618 (93.5)	9002 (100.0)
Revisions	4589 (11.4)	2603 (8.5)	2287 (8.9)	2634 (6.5)	0 (0.0)
Gender					
Males	24962 (62.2)	16013 (60.5)	13179 (56.5)	23037 (57.2)	6481 (72.0)
Females	15147 (37.8)	10444 (39.5)	10158 (43.5)	17215 (42.8)	2521 (28.0)
Age years (at time of surgery)					
<25	17968 (44.8)	11667 (44.1)	10328 (44.3)	20048 (49.8)	Not Reported
≥25	22145 (55.2)	14798 (55.9)	13009 (55.7)	20204 (50.2)	Not Reported

### Outcomes

Reoperations	3762 (10.6)	Not Reported	769 (3.3)	Not Reported	Not Reported
Contralateral Knee Operations	1238 (3.5)	Not Reported	729 (3.1)	1442 (3.8)	Not Reported
Revisions 100 persons-yrs	1209 (3.4)	Not Reported	1105 (4.7)	1601 (4.3)	Not Reported
1 year	0.91	Not Reported	0.84	Not Reported	Not Reported
3 year	1.21	Not Reported	1.19	Not Reported	Not Reported

# Cardiac Device Registry



## Description:

The cardiac device registry was established in 2000, and tracks pacemakers (PM) and implantable cardioverter defibrillators (ICD) and cardiac resynchronization therapy (CRT). As of Q4 2017, there are 128,038 devices in the registry (93,414 initial and 34,624 replacements). The registry volume by device type is noted on page 20.

## Summary:

- **Device Revisions:** The registry tracks all devices undergoing a procedure to explant or replace the device for any reason. Normal battery depletion (ERI) is an expected replacement procedure. Device explant reasons other than ERI include: premature battery depletion, device upgrade/downgrade, mechanical complication of the pulse generator, mechanical complication of a lead, infection, device recall/advisory, pocket erosion/device migration, pocket pain and other patient anatomy issues. Data is available for quality reporting, research, and medical center specific requests. The overall complication rate for devices, excluding normal ERI, implanted from 2007-2017 is noted on page 20.

- **Lead Revisions:** The registry tracks all leads undergoing a procedure to replace, reposition, or repair the lead due to a mechanical malfunction including: lead dislodgement, perforation, conductor fracture, insulation failure, high/low thresholds, oversensing, undersensing, non-capture, extracardiac stimulation, and lead noise. The overall revision rate for leads implanted from 2007-2017 is noted on page 20.
- **Remote Monitoring:** Many cardiac devices have remote monitoring capability which allow the device to transmit data to clinicians without the patient needing to physically come in to the office; therefore, reducing time, effort, and burden on the patient. In collaboration with medical centers in several regions, the registry has identified cardiac device patients who are currently enrolled in remote monitoring and those who still need to be enrolled. Through this feedback mechanism the rates of remote monitoring has increased significantly.

**Registry Champions:** Nigel Gupta, MD, PhD; Kelly Richardson, MD; Cesar Alberte-Lista, MD; Angela Morello, MD; Jason Rashkin, MD; Brant Liu, MD; Jitesh Vasadia, MD; Rasoul Mokabberi, MD

# Cardiac Device Registry

## Registry volume by device type (2007-2017)

Device	# Implants	Initial Volume	Replacement Volume
Pacemaker	86,827	65,508	21,319
ICD	29,824	20,965	8,859
CRT	11,387	6,941	4,446
Total	128,038	93,414	34,624

## Overall complication rate implanted, for devices (excluding normal ERI) (2007-Q2 2017)

Device	Total Volume	Complications	% Complication Rate
<b>Pacemaker</b>			
Dual	48,518	538	1.1
Single	7,962	54	0.67
Leadless	20	1	5
<b>ICD</b>			
Dual	10,516	171	1.62
Single	9,493	80	0.84
<b>CRT</b>			
D	8,201	466	5.68
P	849	25	2.94

## Overall complication rate implanted, for leads (2007-Q2 2017)

Function	Total Volume	Complication Volume	% Complication Rate
BRADY	89,923	1,777	1.97
HF	6,352	201	3.16
TACHY	19,537	614	3.14

# Endovascular Stent Graft Registry



## Description:

This registry was established in 2010 and tracks endovascular stent grafts used for the repair of abdominal aortic aneurysm (AAA). The procedure used to place the stent grafts is endovascular aneurysm repair (EVAR). Currently the registry tracks 3,716 primary and 329 revision procedures.

## Clinical Findings

- Aneurysm size is of clinical importance in evaluating the need for an EVAR procedure. In a total number of 3717 EVAR cases in the registry, the most common aneurysm size for initial EVAR procedures was size 6.0 cm or greater (32.7%) in males, and size 5.0 to 5.9 cm (42.1%) in females. The highest number of reinterventions (16.72%) took place in patients with aneurysm size of 6.0 cm or greater. Registry findings in regard to aneurysm size support regular surveillance and tracking of aneurysm size in patients with known AAAs prior to an EVAR procedure.

- Revisions, secondary interventions, and conversion to open repair are tracked by the registry data. The most common reason for reintervention was endoleak (225 reinterventions out of 3717 total cases). Revisions of the stent graft occurred in 3.61% of all cases (134 revisions out of 3717 total cases). Other reasons for revision can include device malfunction/migration, thrombosis, infection, rupture, etc.
- The majority of patients (58.8%) had a hospital length of stay of 0 to 1 days, with the next highest length of hospital stay being 2 days (18.2% of patients).

## Safety Advisory

- The registry continued to support surgeons in responding to an Endologix Safety Advisory due to higher than anticipated type III endoleaks. By providing KP surgeons and medical centers with a list of patients with affected implants, the registry identified patients at risk, so patients receive appropriate post-market surveillance of their device and treatment as needed.

**Registry Champions:** Bradley Hill, MD; Jeffrey Hsu, MD; Nicolas Nelken, MD; Thomas Rehring, MD; Homayon Hajarizadeh, MD

# Hip Fracture Registry



## Description:

Established in 2009, the hip fracture registry tracks primary surgery of the proximal femur. As of year end 2016, 44,221 primary hip fracture cases and 1,587 revisions were tracked.

## Clinical Findings:

- When compared with Regional Anesthesia (RA), General Anesthesia (GA) is associated with a higher risk of in-hospital mortality, shorter time-to-in-hospital mortality, lower home discharge, and longer time-to-home or health care facility discharge. Also, the previously underrepresented Conversion from Regional to General Anesthesia group demonstrated a higher risk of in-hospital mortality and a shorter time-to-in-hospital mortality when compared with RA. The choice of anesthesia technique can affect in-hospital outcomes, and RA may offer advantages over GA for fragility hip fracture surgeries when possible. (Qiu et al. 2018)
- Use of general anesthesia and conversion from regional to general anesthesia were found to be associated with a higher risk of mortality during the in-hospital stay compared with regional anesthetic techniques, but this higher risk did not persist after hospital discharge. General anesthesia was also found to be associated with a higher risk of all-cause readmission compared with regional, but no other differences were observed in risk for complications. Findings suggest regional anesthetic techniques may be preferred when possible in this patient population. (Desai et al. 2018)
- A study of 39 pediatric patients, < 21 years old at the time of hip fracture, were treated surgically for the hip fractures between 2009 and 2012 in a large integrated health care system. Patient characteristics, type of fracture, surgical treatment, and short-term complications were identified. Hip fractures in a pediatric population were rare, complication rates were low, and few patients had any comorbidities. Hip fractures were more common in boys and Hispanic patients. Intertrochanteric fractures (Delbet Type IV) were the most frequently observed type. (Prentice et al. 2017)

**Registry Champions:** Gary Zohman, MD; Christopher Grimsrud, MD; James Jackman, MD; Kanu Okike, MD

# Hip Fracture Registry

## Hip Fracture KP compared to benchmarks

REGISTRY	Kaiser Permanente 2009-2017	National Hip Fracture Database (NHFD) Annual Report 2017: UK-Wales-Northern Ireland 2007-2016	Swedish Hip Arthroplasty Register 2000-2016	Irish Hip Fracture Database National Report 2014-2016	Australian & New Zealand Hip Fracture Registry (ANZHR)	Norwegian Hip Fracture Register 2005-2016
Data Period	2017	2016	2016	2016	2015-2016	2016
Volume	44,221	500,000	6,158	3,159	12,219	96,597
Female	69%	Not Reported	59%	69%	AUS: 70% / NZ: 68%	70%
Mean Age	Male: 80 Female: 85	Not Reported	Male: 80 Female: 82	Male: 79 Female: 81	AUS: Mean: 82 Median: Male 83, Median: Female: 85 NZ: Mean: 83 Median: Male 85 Median: Female: 85	Mean: Overall: 80 Female: 82 Male: 77
Time to Surgery	Median 21.5 hours 90% within 48 hours	Mean: 32.9 hours Median 24.1 hours 71% within 2 days	Mean: 1.2 days Median: 1 day	75% within 48 hours 59% within 36 hours 41% within 24 hours	AUS: Median 29 hours 82% within 48 hours NZ: Median 24 hours 77% within 48 hours	85% within 48 hours (excluding THAs)
Length of Stay	Mean: 4.8 days	Mean: 16.6 days Median: 12 days	Mean: 10.7 days Median: 9 days	Mean: 20days Median 12days	AUS: Median 7.7 days NZ: Median 5.8 days	Mean: 9.8 days
Revision Rate	3.59%	Not Reported	3.30%	5% within 30days	Not Reported	4.00%
Mortality	11.0%	6.7%	13.0%	Not Reported	5% before discharge, additional 15-20% within 1 year	Not Reported

# Shoulder Arthroplasty Registry



## Description:

Kaiser Permanente tracks elective and urgent shoulder arthroplasty procedures including total shoulder arthroplasty (TSA), reverse total shoulder arthroplasty (RTSA), hemiarthroplasty (HA) and humeral head resurfacing (HHR) procedures. Between 2005 and year end 2016, the shoulder registry has captured information for 14,301 primary procedures.

## Clinical Findings

- In 5,121 Shoulder Arthroplasties 75.6% of patients were opioid users 1-year pre-operatively: 92.5% were opioid users in Q1 post-operatively, 40-44% were opioid users in Q2-Q4 post-operatively. Risk factors found to be associated with increased opioid utilization 1-year postoperatively: Preoperative opioid utilization, age >60, opioid dependence, anxiety, history of substance abuse, and general chronic pain. These findings emphasize the need for surgeon and patient awareness, and education in the management of postoperative opioid usage associated with the indicated conditions.

- In 5,487 Shoulder Arthroplasties, infection risk was significantly lower in TSA procedures using antibiotic-laden irrigation but not in RTSA or HA. There was no association of infection prevention with antibiotic cement in TSA and RTSA. Spacesuit use was not associated with infection risk in TSA, RTSA, or HA. Further prospective investigations are needed to identify safe and effective ancillary antibiotic administration practices.
- In 7,421 primary Shoulder Arthroplasties, diabetes was not associated with short-term revision, deep infection, VTE, or rehospitalization after shoulder arthroplasty. No significant association was found between these complications and glycemic control, as measured by preoperative and postoperative HbA1c, in diabetic patients. The strict application of thresholds of acceptable preoperative HbA1c levels may not be an effective method of reducing infection rates and other possible diabetes associated complications in shoulder arthroplasty.

**Registry Champions:** Ronald Navarro, MD; Mark Dillon, MD; Mark Shaieb, MD; Terrill Julien, MD; Darin Allred, MD; Matthew Budge, MD; Anita Rao, MD



# Shoulder Arthroplasty Registry

Shoulder Arthroplasty KP compared to benchmarks			
	Kaiser Permanente	Australia Orthopaedic Association Shoulder Arthroplasty	The New Zealand Joint Registry
Time Period	2005-2016	2004-2016	2000-2016
Volume	14301	38265	8250
Gender, Female	55.12%	62.10%	63.14%
Mean Age (yrs)	69.61	71.4	71.11
Revision Rate	TSA: 0.70/ 100 obs yrs RTSA: 1.21/ 100 obs yrs	1.70/ 100 obs yrs 1.23/ 100 obs yrs	0.99/ 100 obs yrs
Top 3 Reasons for Primary	Osteoarthritis Rotator Cuff Arthropathy Fracture	Osteoarthritis Rotator Cuff Arthropathy Fracture	Osteoarthritis Cuff Tear Arthropathy Fracture of Proximal Humerus
Top 3 Reasons for Revision	Instability/Dislocation Infection Rotator Cuff Tear	Instability/Dislocation Loosening Rotator Cuff Insufficiency	Pain Instability/Dislocation Subacromial Cuff Impingement
Outcome			
Infection	0.90%	Not Reported	Not Reported
DVT	0.65%	Not Reported	Not Reported
PE	0.52%	Not Reported	Not Reported

# Spine Registry



## Description

Implemented in 2009, this registry tracks over 23,790 instrumented and non-instrumented spinal procedures performed by the Neurosurgery and Orthopedic Spine surgeons as of year-end 2016. This represents more than 80,000 total implants.

## Clinical Findings

- Every 5 kg/m<sup>2</sup> increase in BMI is associated with a significant increase in surgical time (7.8 minutes), estimated blood loss (EBL) (36.5 mL), risk of deep infection (OR =1.7 times), and deep vein thrombosis (DVT) (OR =1.5). BMI was not associated with increased incidence of other intraoperative or health-related complications. Rate of re-operation was 1.1 times higher with every 5 kg/m<sup>2</sup> increase in BMI, but rate of re-operation due to adjacent segment disease (ASD) was not associated with BMI in our spine patient population. (Akins, et al. 2017)


- In 10,416 patients our preliminary study found no evidence that exogenous, local BMP administration during spine fusion influences subsequent cancer risk. Additional studies and longer follow-up is necessary to definitively conclude our findings. (Bains, et al. 2017)
- In a cohort of 1158 posterior cervical spine fusion patients reoperation rates for symptomatic nonunions in subaxial spine (C2-7) with more than 1 year of follow-up were found to be 1.1% with BMP and 0.7% without BMP. There was no significant difference in the reoperation rates for symptomatic nonunions with or without BMP. (Guppy, et al. 2016)

**Registry Champions:** Kern Guppy, PhD, MD; Johannes Bernbeck, MD; Harsimran Brara, MD

# Spine Surgery Registry

<b>Spine Surgery KP compared to benchmarks</b>		
<b>Variable</b>	<b>Kaiser Permanente</b>	<b>EUROSPINE</b>
Time Period	2009-2013, 2016	2006-2016
Volume	23,790	102,025
<b>Demographics</b>		
Age (Mean)	57.7 years	56.8
Gender	51.9% female	50.9% female
Current Smoker	7.6%	Not Reported
Diagnosis - Degenerative	70.6%	80.3%
<b>Fusion Approach</b>		
Anterior Only	24.5%	Not Reported
Posterior Only	63.5%	Not Reported
Combined	12%	Not Reported
<b>Outcomes</b>		
Dural Tear	3.2%	4.8%
Superficial Infection	0.6%	2.2%
Deep Infection	0.6%	4.4%
Nonunion	1.6%	20.9%
Adjacent Segment Disease	4.4%	25.6%

# Total Joint Replacement Registry



## Description

The Total Joint Replacement Registry (TJRR), established in 2001, collects patient and surgical implant information and patient outcomes. The TJRR now tracks over 250,000 procedures through December 2016 (242,469 primary total joint replacements and 6,992 revision cases).

## Clinical findings

- In 47,523 primary THA cases, 95.2% and 4.8% had Osteoarthritis and Osteonecrosis, respectively. The osteonecrosis cohort had higher crude incidence of 90-day mortality, SSI, unplanned readmission and revision. Compared to OA, a diagnosis of osteonecrosis was associated with worse outcomes post-THA. A detailed preoperative discussion including the risk of complications is needed for informed consent from patients with osteonecrosis. (Sindh et al. 2017)
- In a study evaluating risk of all-cause revision in metal versus ceramic femoral heads when used with an highly cross-linked polyethylene liner, including an evaluation of the effect of head size; and dislocation in metal versus ceramic femoral heads when used

with an HXLPE liner, for all-cause revision, there was no difference between ceramic versus metal heads in combination with an HXLPE liner. Smaller metal head sizes of <32 mm were associated with increased risk of revision relative to 36 mm. For dislocation, ceramic heads increased risk relative to metal at < 32 mm only. Head sizes < 32 mm were associated with increased risk of dislocation relative to 36 mm for metal and ceramic heads. The results did not provide evidence for use of one femoral head material over another when used with HXLPE liners for the outcome of revision, but for dislocation, metal performed better than ceramic with < 32-mm heads. Overall, the findings suggest increased risk of revision/dislocation with head sizes < 32 mm. (Cafri et al. 2017)

- Based on study results, fondaparinux, enoxaparin, and warfarin was not found to differ from aspirin in the prevention of PE, DVT, or VTE. Furthermore, aspirin was not found to differ from the alternative chemoprophylactics with respect to safety endpoints. When specifically testing for noninferiority, enoxaparin was found to be as safe as aspirin with respect to bleeding and fondaparinux as safe as aspirin for risk of wound complications. (Cafri et al 2017)

**Registry Champions:** Mark Melberg MD; Le-Don A Robinson MD; Gregory Y Lee MD; Maurice Cates MD; Adrian D Hinman MD; Erik W Kroger MD; Matthew P Kelly MD; Thomas Stoll MD

# Total Joint Replacement Registry

## Total Hip Replacement KP compared to benchmarks


Registry	Data	Total Cases	Revision % Period	Age	Female %	10-yr. Survival % (CI)
Kaiser Permanente	2001-2016	86,376	3.1%	65.5 mean age	50.8%	95.1 (94.8-95.3)
Australia	1999-2015	346,782	4.3%	67.7 mean age	55.1%	93.4 (93.2-93.5)
United Kingdom	2003-2016	890,681	2.7%	69 median age (61-76 IQR)	59.8%	94.79 (94.71-94.87)
Sweden	1991-2015	163,341	5.6%	Mean age: Male 67.3, Female 70.0	60.0%	94.7 (94.4-94.9)
Norway	1987-2016	181,048	8.7%	Mean age: Male 67.1, Female 69.9	67.1%	91.5 (91.3-91.7)

## Total Knee Replacement KP compared to benchmarks

Registry	Data	Total Cases	Revision % Period	Age	Female %	10-yr. Survival % (CI)
Kaiser Permanente	2001-2016	163,093	2.7%	67.3 mean age	54.7%	95.7 (95.5-95.8)
Australia	1999-2015	494,571	9.8%	68.6 mean age	56.9%	94.7 (94.6-94.8)
United Kingdom	2003-2016	975,739	5.6%	69 median age (63-76 IQR)	56.8%	96.61 (96.55-96.67)
Sweden	1999-2015	163,120	4.3%	68.6 mean age	59.7%	96 (95.9-96.1)
Norway	1994-2016	75,229	5.9%	Mean age: Male 67.4, Female 69.2	63.5%	92.2 (92-92.4)


# Total Joint Replacement Registry

## Annual Total Hip Arthroplasty (THA) Revision Burden



	2011	2012	2013	2014	2015	2016
■ KP	8.2%	10.0%	9.1%	8.7%	2.9%	3.1%
■ AJRR	—	14.6%	14.9%	13.9%	12.3%	8.6%
■ NJR	10.9%	11.7%	10.4%	9.0%	2.6%	2.7%
■ NAR	14.9%	14.3%	13.9%	13.7%	14.2%	14.2%

## Annual Total Knee Arthroplasty (TKA) Revision Burden



	2011	2012	2013	2014	2015	2016
■ KP	5.3%	5.9%	5.7%	5.7%	2.6%	2.7%
■ AJRR	—	6.2%	6.1%	7.3%	6.3%	5.1%
■ NJR	5.6%	6.1%	5.6%	5.2%	2.3%	5.6%
■ NAR	8.2%	8.5%	7.9%	7.5%	8.4%	9.1%

### Registries:

KP: Kaiser Permanente Implant Registries  
 AJRR: American Joint Replacement Registry  
 NJR: National Joint Registry, UK  
 NAR: Norwegian Arthroplasty Register

# Implant Registries Staff

## Management Team

Liz Paxton, MA	Director
Jessica Harris, MS, RD	Manager
Kenneth Sucher, MS	Manager

## Administrative Team

Betty Jo Alfstad, MS	Staff Specialist
Donna Leck	Research Administrative Analyst

## Biostatisticians/Research Scientists

Guy Cafri, PhD, MStat	Research Scientist Biostatistician
Priscilla Chan, MS,	Data Consultant
Heather Prentice, PhD, MPH	Research Scientist Investigator
Wei Wang, PhD, MS	Biostatistician III

## Database Administrators

Kim Phan, BA	Senior Application Development Technical Specialist
Buu Truong, BS, MIS	Senior Application Program Analyst

## Data Consultants/Programmers

Yuexin (Cindy) Chen, BS	Senior Data Consultant
Tomy Huon, BS	SAS Programmer III
Michael Reyes, BS	SAS Programmer II
Craig Salman, BS	SAS Programmer II
Xueyan (Sophie) Su, BS	SAS Programmer II
Hong (Maggie) Sun, MS	SAS Programmer II

## Data Quality

Jalessia Moore	Research Associate I
Chelsea Reyes	Project Analyst
Nicole Sankowski, BS	Research Associate I

## Project Managers

William Burfeind, MAS	Project Manager III
Brian Fasig, PhD	Project Manager III

## Quality Reporting

Anna Ritko, MA, MPhil	Senior Data Consultant
-----------------------	------------------------

## Research Associates

Janine Cruz, BA	Research Associate II
Kristen Mays, MA	Research Associate III
Tia Mullane, BA	Research Associate II
Juan Ruiz	Research Associate II
Andrew (Scott) Thomas, BS	Research Associate II

# Publications

## Anterior Cruciate Ligament Reconstruction

### 2017

Arianjam A, Inacio MCS, Funahashi TT, Maletis GB. Analysis of 2019 patients undergoing revision anterior cruciate ligament reconstruction from a community-based registry. *Am J Sports Med*, 2017; 45(7): 1574-80.

Davis BR, Chen J, Inacio MCS, Love R, Prentice HA, Maletis GB. The incidence of subsequent meniscal surgery is higher in the anterior cruciate ligament-reconstructed knee than in the contralateral knee. *Am J Sports Med*, 2017; 45(14): 3216-22.

Maletis GB, Chen J, Inacio MCS, Love RM, Funahashi TT. Increased risk of revision after anterior cruciate ligament reconstruction with bone-patellar tendon-bone allografts compared with autografts. *Am J Sports Med*, 2017; 45(6): 1333-40.

Maletis GB, Chen J, Inacio MCS, Love RM, Funahashi TT. Increased risk of revision after anterior cruciate ligament reconstruction with soft tissue allografts compared with autografts. *Am J Sports Med*, 2017; 45(8): 1837-44.

Wyatt RWB, Inacio MCS, Bellevue KD, Schepps AL, Maletis GB. Isolated ACL versus multiple knee ligament injury: Associations with patient characteristics, cartilage status, and meniscal tears identified during ACL reconstruction. *Phys Sportsmed*, 2017; 45(3): 323-8.

Yu A, Prentice HA, Burfeind WE, Funahashi T, Maletis GB. Risk of infection after allograft anterior cruciate ligament reconstruction: Are nonprocessed allografts more likely to get infected? *Am J Sports Med*, 2018; Epub ahead of print.

### 2016

Nelson I, Chen J, Love R, Davis B, Maletis G, Funahashi T. (2016). A comparison of revision and rerupture rates of ACL reconstruction between autografts and allografts in the skeletally immature Knee Surg Sports Traumatol Arthrosc, 24 (3): 773-779.

Spragg L, Chen J, Mirzayan R, Love R, Maletis G. (2016). The Effect of Autologous Hamstring Graft Diameter on the Likelihood for Revision of Anterior Cruciate Ligament Reconstruction *Am J Sports Med*, 44 (6): 1475-1481.

Inacio MC, Cafri G, Funahashi T, Maletis G, Paxton E. (2016). Type and frequency of healthcare encounters can predict poor surgical outcomes in anterior cruciate ligament reconstruction patients *International Journal of Medical Informatics*, 90 32-39.

Tibor L, Chan PH, Funahashi T, Wyatt R, Maletis GB, Inacio MC. (2016). Surgical Technique Trends in Primary ACL Reconstruction from 2007 to 2014 *J Bone Joint Surg Am*, 13 (98): 1079-89.

### 2015

Inacio MC, Chen Y, Maletis GB, Ake CF, Fithian DC, Granan LP. (2015). Injury pathology at the time of anterior cruciate ligament reconstruction associations with self-assessment of knee function. *Clin J Sport Med*, 24(6): 461-467.

Maletis GB, Chen J, Inacio MC, Fundahashi TT. (2015). Age-Related Risk Factors for Revision Anterior Cruciate Ligament Reconstruction: A Cohort Study of 21,304 Patients From the Kaiser Permanente Anterior Cruciate Ligament Registry. *Am J Sports Med*, 44(2): 331-336.

Navarro RA, Inacio MC, Maletis GB. (2015). Does Racial Variation Influence Preoperative Characteristics and Intraoperative Findings in Patients Undergoing Anterior Cruciate Ligament Reconstruction? *Am J Sports Med*, 43(12): 2959-2965.

Tejwani SG, Chen J, Funahashi TT, Maletis GB, Love R. (2015). Revision Risk After Allograft Anterior Cruciate Ligament Reconstruction: Association With Graft Processing Techniques, Patient Characteristics, and Graft Type. *Am J Sports Med*, 43(11): 2696-2705.

### 2014

Bojicic JL, Sue VM, Huon TS, Maletis GB, Inacio MC. (2014). Comparison of paper and electronic surveys for measuring patient-reported outcomes after anterior cruciate ligament reconstruction. *Perm J*, 18(3): 22-6.



# Publications

## Anterior Cruciate Ligament Reconstruction

*continued from page 32*

Funahashi KM, Moksnes H, Maletis GB, Csintalan RP, Inacio MC, Funahashi TT. (2014). Anterior Cruciate Ligament Injuries in Adolescents With Open Physis: Effect Of Recurrent Injury and Surgical Delay On Meniscal and Cartilage Injuries. *Am J Sports Med*, 42(5): 1068-73.

Granán LP, Baste V, Engebretsen L, Inacio MC. (2014). Associations Between Inadequate Knee Function Detected By KOOS and Prospective Graft Failure in an Anterior Cruciate Ligament-Reconstructed Knee. *Knee Surg Sports Traumatol Arthrosc*, 23(4): 1135-1140.

Maletis GB, Inacio MC, Funahashi TT. (2014). Risk Factors Associated With Revision and Contralateral Anterior Cruciate Ligament Reconstructions in the Kaiser Permanente ACLR Registry. *Am J Sports Med*, 43(3): 641-647.

Wyatt RW, Inacio MC, Liddle KD, Maletis GB. (2014). Prevalence and Incidence of Cartilage Injuries and Meniscus Tears in Patients Who Underwent Both Primary and Revision Anterior Cruciate Ligament Reconstructions. *Am J Sports Med*, 42(8): 1841-1846.

### 2013

Csintalan R, Inacio MC, Funahashi T, Maletis GB. (2013). Risk Factors of Subsequent Operations After Primary Anterior Cruciate Ligament Reconstruction. *Am J Sports Med*, 42(3): 619-625.

Granán L, Inacio M, Maletis G, Funahashi T, Engebretsen L. (2013). Sport Specific Injury Pattern Recorded During Anterior Cruciate Ligament Reconstruction. *Am J Sports Med*, 41(12): 2814-2818.

Maletis GB, Inacio M, Reynolds S, Desmond J, Maletis M, Funahashi T. (2013). Incidence of Postoperative Anterior Cruciate Ligament Reconstruction Infections: Graft Choice Makes a Difference. *Am J Sports Med*, 41(8): 1780-5.

Maletis GB, Inacio MC, Desmond JL, Funahashi

TT. (2013). Reconstruction of the Anterior Cruciate Ligament: Association of Graft Choice with Increased Risk of Early Revision. *Bone Joint J*, 95-B(5): 623-8.

Maletis GB, Inacio MC, Funahashi TT. (2013). Analysis of 16,192 Anterior Cruciate Ligament Reconstructions From a Community-Based Registry. *Am J Sports Med*, 41(9): 2090-8.

Paxton E, Inacio M, Maletis G, Desmond J, Funahashi T. (2013). What Have We Learned from the Kaiser Permanente Anterior Cruciate Ligament Reconstruction Registry (KP ACLRR). *The ACL-Deficient Knee*, 3-10(Springer-Verlag): 3-10.

Wyatt RW, Inacio MC, Liddle KD, Maletis GB. (2013). Factors Associated With Meniscus Repair in Patients Undergoing Anterior Cruciate Ligament Reconstruction. *Am J Sports Med*, 41(12): 2766-71.

### 2012

Csintalan RP, Inacio MCS, Desmond JL, Funahashi TT. (2012). Anterior Cruciate Ligament Reconstruction in Patients with Open Physes: Early Outcomes. *J Knee Surg*, 26(4): 225-32.

Granán LP, Inacio MC, Maletis GB, Funahashi TT, Engebretsen L. (2012). Intraoperative Findings and Procedures in Culturally and Geographically Different Patient and Surgeon Populations. *Acta Orthopaedica*, 83(6): 577-82.

Inacio M, Maletis G, Paxton E, Granán L, Csintalan R, Fithian D, Funahashi T. (2012). Patient and Surgeon Characteristics Associated with Primary Anterior Cruciate Ligament Reconstruction Graft Selection. *Am J Sports Med*, 40(2): 339-45.

### 2011

Chhadia A, Inacio M, Maletis G, Csintalan R, Davis B, Funahashi T. (2011). Are Meniscus and Cartilage Injury Related to Time to Anterior Cruciate Ligament Reconstruction? *Am J Sports Med*, 39(9): 1894-9.

# Publications

## Anterior Cruciate Ligament Reconstruction

*continued from page 33*

Maletis G, Granan LP, Inacio M, Funahashi T, Engebretsen L. (2011). Comparison of a Community Based Anterior Cruciate Ligament Reconstruction Registry in the United States and Norway. *J Bone Joint Surg Am*, 93(Supplement 3): 31-6.

### 2009

Mehta VM, Paxton EW, Fithian DC. (2009). Does the Use of Fluoroscopy and Isometry During Anterior Cruciate Ligament Reconstruction Affect Surgical Decision Making? *Clin J Sport Med*, 19(1): 46-8.

### 2008

Csintalan RP, Inacio MC, Funahashi TT, Csintalan RP, Inacio MCS, Funahashi TT. (2008). Incidence Rate of Anterior Cruciate Ligament Reconstructions. *Perm J*, 12(3): 17-21.

Mehta VM, Young EP, Paxton EW, Fithian DC. (2008). The Effect of Ketorolac on Anteroposterior Knee Laxity After Anterior Cruciate Ligament Reconstruction. *Orthopedics*, 31(6): 538-40.

### 2007

Mehta VM, Paxton LW, Fornalski SX, Csintalan RP, Fithian DC. (2007). Reliability of the International Knee Documentation Committee Radiographic Grading System. *Am J Sports Med*, 35(6): 933-5.

### 2005

Crain EH, Fithian DC, Paxton EW, Luetzow WF. (2005). Variation in Anterior Cruciate Ligament Scar Pattern: Does The Scar Pattern Affect Anterior Laxity in Anterior Cruciate Ligament-Deficient Knees? *J Arthroplasty*, 21(1): 19-24.

Fithian DC, Paxton EW, Stone ML, Luetzow WF, Csintalan RP, Phelan D, Daniel DM. (2005). Prospective Trial of a Treatment Algorithm for the Management of the Anterior Cruciate Ligament-Injured Knee. *Am J Sports Med*, 33(3): 335-46.

Theodorou D TS, Fithian D, Paxton E, Garelick D, Resnick D. (2005). Posterolateral Complex Knee Injuries: MRI with Surgical Correlation. *Acta Radiol*, 46(3): 297-305.

### 2003

Ohara W, Paxton E, Fithian C. (2003). Epidemiology of Knee Ligament Injuries. *Pedowitz, Robert A., O'Connor, John J., and Akeson, Wayne H.. Daniel's Knee Injuries : Ligament and Cartilage Structure, Function, Injury, and Repair(2nd edition)*: 311- 344.

Paxton EW, Fithian DC, Stone ML, Silva P. (2003). The reliability and validity of knee-specific and general health instruments in assessing acute patellar dislocation outcomes. *Am J Sports Med*, 31(4): 487-92.

### 2002

Fithian D, Paxton E, Goltz D. (2002). Fate of the anterior cruciate ligament-injured knee. *Orthop Clin North Am*, 33(4): 621- 636.

## Cardiac Device

### 2015

Krishnaswami A, Kiley M, Anthony F, Chen Y, Chen J, Rajagopal S, Liu T, Young C, Paxton E. (2015). Effect of Advancing Age and Multiple Chronic Conditions on Mortality in Patients with End-Stage Renal Disease after Implantable Cardioverter- Defibrillator Placement. *Perm J*, 20(1): 27-32.

## Endovascular Stents

### 2016

Gupta N, Kiley ML, Anthony F, Young C, Brar S. (2016). Multi-Center, Community-Based Cardiac Implantable Electronic Devices Registry: Population, Device Utilization, and Outcomes *J Am Heart Assoc*, 4 (3): 1-9.

# Publications

## Endovascular Stents

*continued from page 34*

### 2015

Hye RJ, Inui TS, Anthony FF, Kiley ML, Chang RW, Rehling TF, Nelken NA, Hill BB. (2015). A Multiregional Registry Experience Using an Electronic Medical Record to Optimize Data Capture for Longitudinal Outcomes in Endovascular Abdominal Aortic Aneurysm Repair. *J Vasc Surg*, 61(5): 1160-1167.

## Heart Valve Replacement

### 2013

Chen JC, Pfeffer T, Johnstone S, Chen Y, Kiley M-L, Richter R, Lee H. (2013). Analysis of Mitral Valve Replacement Outcomes is Enhanced by Meaningful Clinical Use of Electronic Health Records. *Perm J*, 17(2): 12-6.

## Hip Fracture

### 2017

Prentice HA, Paxton EW, Hunt JJ, Grimsrud CD, Weiss JM. "Pediatric hip fractures in California: Results from a community-based hip fracture registry." *Perm J*, 2017; 21: 16-081.

Okike K, Chan PH, Paxton EW. "Effect of surgeon and hospital volume on morbidity and mortality after hip fracture." *J Bone Joint Surg Am*, 2017; 99(18): 1547-53.

### 2016

Brox T, Chan PH, Cafri G, Inacio MC. (2016). Similar mortality with general or regional anesthesia in elderly hip fracture patients *Acta Orthopaedica*, 87 (2): 152-157.

### 2015

Inacio MC, Weiss JM, Miric A, Hunt JJ, Zohman GL, Paxton EW. (2015). Community-Based Hip Fracture Registry: Population, Methods, and Outcomes. *Perm J*, 19(3): 29-36.

## Shoulder Arthroplasty Replacement

### 2017

Anakwenze O, Fokin A, Chocas M, Dillon MT, Navarro RA, Yian EH, Singh A. Complications in total shoulder and reverse total shoulder arthroplasty by body mass index. *J Shoulder Elbow Surg*, 2017; 26(7): 1230-7.

Anakwenze O, Yehyawi T, Dillon MT, Paxton E, Navarro RA, Singh A. Effect of age on outcomes of shoulder arthroplasty. *Perm J*, 2017; 21: 16-056.

Dillon MT, Chan PH, Inacio MCS, Singh A, Yian EH, Navarro RA. Yearly trends in elective shoulder arthroplasty, 2005–2013. *Arthritis Care (Hoboken)*, 2017; 69(10): 1574-81.

Paxton EW, Prentice HA, Inacio MCS, Dillon MT, Page RS, Rasmussen JV, Salomonsson B, de Steiger R, Navarro RA. Are we throwing the baby out with the bath water? *J Shoulder Elbow Surg*, 2017; 26(5): e137-9.

### 2016

Singh A, Padilla M, Nyberg E, Chocas M, Anakwenze O, Mirzayan R, Yian E, Navarro RA. (2016). Cement technique correlates with tuberosity healing in hemiarthroplasty for proximal humeral fracture *J Shoulder Elbow Surg*, 26 (3): 437-442.

### 2015

Dillon MT, Ake CF, Burke MF, Singh A, Yian EH, Paxton EW, Navarro RA. (2015). The Kaiser Permanente Shoulder Arthroplasty Registry. *Acta Orthopaedica*, 86(3): 286-292.

# Publications

## Shoulder Arthroplasty Replacement

*continued from page 35*

Yian EH, Dillon M, Sodl J, Dionysian E, Navarro R, Singh A. (2015). Incidence of symptomatic compressive peripheral neuropathy after shoulder replacement. *Hand*, 10(2): 243-247.

### 2014

Inacio MC, Dillon MT, Miric A, Anthony FH, Navarro RA, Paxton EW. (2014). Mortality After Shoulder Arthroplasty. *J Arthroplasty*, 29(9): 1823–1826.

Page RS, Navarro RA, Salomonsson B. (2014). Establishing an International Shoulder Arthroplasty Consortium. *J Shoulder Elbow Surg*, 23(8): 1081-2.

Richards J, Inacio MC, Beckett M, Navarro RA, Singh A, Dillon MT, Sodl JF, Yian EH. (2014). Patient and procedure-specific risk factors for deep infection after primary shoulder arthroplasty. *Clin Orthop Relat Res*, 472(9): 2809-15.

Singh A, Yian E, Dillon M, Burke M, Takayanagi M, Navarro R. (2014). The Effect of Surgical Volume on Shoulder Arthroplasty Quality Metrics. *J Shoulder Elbow Surg*, 23(8): 1187-94.

### 2013

Dillon MT, Inacio MC, Burke MF, Navarro RA, Yian EH. (2013). Shoulder Arthroplasty in Patients 59 Years of Age and Younger. *J Shoulder Elbow Surg*, 22(10): 1338-44.

Navarro RA, Inacio MC, Burke MF, Costouros JG, Yian EH. (2013). Risk of Thromboembolism in Shoulder Arthroplasty: Effect of Implant Type and Traumatic Indication. *Clin Orthop Relat Res*, 471(5): 1576-81.

### 2007

Sachs RA, Lin D, Stone ML, Paxton E, Kuney M. (2007). Can the Need for Future Surgery for Acute Traumatic Anterior Shoulder Dislocation Be Predicted? *J Bone Joint Surg Am*, 89(8): 1665-74.

### 2005

Sachs RA, Williams B, Stone ML, Paxton L, Kuney M. (2005). Open Bankart Repair: Correlation of Results with Postoperative Subscapularis Function. *Am J Sports Med*, 33(10): 1458-62.

### 2003

Green LB, Pietrobon R, Paxton E, Higgins LD, Fithian D. (2003). Sources of Variation in Readmission Rates, Length of Stay, and Operative Time Associated with Rotator Cuff Surgery. *J Bone Joint Surg Am*, 85-A(9): 1784-9.

## Spine

### 2017

Akins PT, Inacio MCS, Bernbeck JA, Harris J, Chen YX, Prentice HA, Guppy KH. Do obese and extremely obese patients lose weight after lumbar spine fusions? Analysis of a cohort of 7303 patients from the Kaiser National Spine Registry. *Spine (Phila Pa 1976)*, 2018; 43(1): 22-7.

Bains R, Mitsunaga L, Kardile M, Chen Y, Guppy K, Harris J, Paxton E. Bone morphogenetic protein (BMP-2) usage and cancer correlation: An analysis of 10,416 spine fusion patients from a multi-center spine registry. *J Clin Neurosci*, 2017; 43: 214-9.

Bains RS, Kardile M, Mitsunaga L, Chen Y, Harris J, Paxton E, Majid K. Does chronic kidney disease affect the mortality rate in patients undergoing spine surgery? *J Clin Neurosci*, 2017; 43: 208-13.

Flippin M, Harris J, Paxton EW, Prentice HA, Fithian DC, Ward SR, Gombatto SP. Effect of body mass index on patient outcomes of surgical intervention for the lumbar spine. *J Spine Surg*, 2017; 3(3): 349-57.

# Publications

## Spine

*continued from page 36*

### 2016

Guppy KH, Harris J, Chen J, Paxton EW, Alvarez JL, Bernbeck JA. (2016). Reoperation rates for symptomatic nonunions in posterior cervical (subaxial) fusions with and without bone morphogenetic protein in a cohort of 1158 patients *J Neurosurg Spine*, 24 (4): 556-564.

Guppy KH, Harris J, Chen J, Paxton EW, Bernbeck JA. (2016). Reoperation rates for symptomatic nonunions in posterior cervicothoracic fusions with and without bone morphogenetic protein in a cohort of 450 patients *J Neurosurg Spine*, 25 (3): 309-317.

### 2015

Akins P, Harris J, Alvarez J, Chen Y, Paxton E, Bernbeck J, Guppy K. (2015). Risk Factors Associated With 30-day Readmissions After Instrumented Spine Surgery in 14,939 Patients: 30-day Readmissions After Instrumented Spine Surgery. *Spine (Phila Pa 1976)*, 40(13): 1022–1032.

Guppy KH, Brara HS, Bernbeck JA. (2015). Occipitocervical Fusions in Elderly Patients: Mortality and Reoperation Rates from a National Spine Registry. *World Neurosurgery*, 86 161–167.

Guppy KH, Harris J, Chen J, Paxton EW, Alvarez JL, Bernbeck JA. (2015). Reoperation rates for symptomatic nonunions in posterior cervical (subaxial) fusions with and without bone morphogenetic protein in a cohort of 1158 patients. *J Neurosurg Spine*, 24(4): 556-564.

Guppy KH, Harris J, Paxton LW, Alvarez JL, Bernbeck JA. (2015). Reoperation Rates for Symptomatic Nonunions in Anterior Cervical Fusions from a National Spine Registry. *Spine*, 40(20): 1632–1637.

### 2014

Guppy K, Paxton E, Harris J, Alvarez J, Bernbeck J. (2014). Does Bone Morphogenetic Protein Change the Operative Nonunion Rates in Spine Fusions? *Spine (Phila Pa 1976)*, 39(22): 1831-1839.

## Total Joint Replacement

### 2017

Bini SA, Cafri G, Khatod M. Midterm-adjusted survival comparing the best performing unicompartmental and total knee arthroplasties in a registry. *J Arthroplasty*, 2017; 32(2017): 3352-5.

Cafri G, Paxton EW, Chen Y, Cheetham CT, Gould MK, Sluggett J, Bini SA, Khatod M. Comparative effectiveness and safety of drug prophylaxis for prevention of venous thromboembolism after total knee arthroplasty. *J Arthroplasty*, 2017; 32(11): 3524-8.

Cafri G, Paxton EW, Love R, Bini SA, Kurtz SM. Is there a difference in revision risk between metal and ceramic heads on highly crosslinked polyethylene liners? *Clin Orthop Relat Res*, 2017; 475(5): 1349-55.

Inacio MCS, Dillon MT, Miric A, Navarro RA, Paxton EW. Mortality after total knee and total hip arthroplasty in a large integrated health care system. *Perm J*, 2017; 21: 16-171.

Inacio MCS, Paxton EW, Graves SE, Namba RS, Nemes S. Projected increase in total knee arthroplasty in the United States – an alternative projection model. *Osteoarthritis Cartilage*, 2017; 25(11): 1797-803.

Namba RS, Inacio MCS, Pratt NL, Graves SE, Roughead EE, Paxton EW. Persistent opioid use following total knee arthroplasty: A signal for close surveillance. *J Arthroplasty*, 2018; 33(2): 331-6.

Singh JA, Chen J, Inacio MCS, Namba RS, Paxton EW. An underlying diagnosis of osteonecrosis of bone is associated with worse outcomes than osteoarthritis after total hip arthroplasty. *BMC Musculoskelet Disord*, 2017; 18(1): 8.

# Publications

## Total Joint Replacement

*continued from page 37*

### 2016

Namba RS, Inacio MC, Pratt NL, Graves SE, Roughead EE, Cheetham CT, Paxton EW. (2016). Postoperative opioid use as an early indication of total hip arthroplasty failure *Acta Orthopaedica*, 362 (87): 37-43.

Paxton EW, Torres A, Love RM, Barber TC, Sheth DS, Inacio MC. (2016). Total joint replacement: A multiple risk factor analysis of physical activity level 1-2 years postoperatively *Acta Orthopaedica* (87): 44-9.

Sheth DS, Cafri G, Paxton EW, Namba RS. (2016). Bilateral Simultaneous vs Staged Total Knee Arthroplasty: A Comparison of Complications and Mortality *J Arthroplasty*, 31 (9): 212-216.

Inacio MC, Pratt NL, Roughead EE, Paxton EW, Graves SE. (2016). Opioid use after total hip arthroplasty surgery is associated with revision surgery *BMC Musculoskeletal Disorders*, 17:122 1-9.

### 2015

Bini SA, Chan PH, Inacio MC, Paxton EW, Khatod M. (2015). Antibiotic cement was associated with half the risk of re-revision in 1,154 aseptic revision total knee arthroplasties. *Acta Orthopaedica*, 87(1): 55-59.

Khatod M, Cafri G, Inacio M, Schepps A, Paxton EW, Bini S. (2015). Revision Total Hip Arthroplasty: Factors Associated with Re-Revision Surgery. *J Bone Joint Surg Am*, 97(5): 359-366.

Khatod M, Inacio MC, Dell RM, Bini SA, Paxton EW, Namba RS. (2015). Association of Bisphosphonate Use and Risk of Revision After THA: Outcomes From a US Total Joint Replacement Registry. *Clin Orthop Relat Res*, 473(11): 3412-3420.

Miric A, Inacio MC, Kelly MP, Namba RS. (2015). Are Nonagenarians Too Old For Total Hip Arthroplasty? An Evaluation of Morbidity and Mortality Within a Total Joint Replacement Registry. *J Arthroplasty*, 30(8): 1324-1327.

Namba RS, Inacio MC, Cheetham TC, Paxton EW, Khatod M. (2015). Lower Total Knee Arthroplasty Revision Risk Associated with Bisphosphonate Use, Even in Patients with Normal Bone Density. *J Arthroplasty*, 31(2): 537-541.

Paxton EW, Inacio MC, Khatod M, Yue E, Funahashi T, Barber T. (2015). Risk Calculators Predict Failures of Knee and Hip Arthroplasties: Findings from a Large Health Maintenance Organization. *Clin Orthop Relat Res*, 473(12): 3965-3973.

Paxton EW, Inacio MC, Kurtz S, Love R, Cafri G, Namba RS. (2015). Is There a Difference in Total Knee Arthroplasty Risk of Revision in Highly Crosslinked versus Conventional Polyethylene? *Clin Orthop Relat Res*, 473(3): 999-1008.

Paxton EW, Inacio MC, Namba RS, Love R, Kurtz SM. (2015). Metal-on-conventional Polyethylene Total Hip Arthroplasty Bearing Surfaces Have a Higher Risk of Revision Than Metal-on-highly Crosslinked Polyethylene: Results From a US Registry. *Clin Orthop Relat Res*, 473(3): 1011-1021.

Paxton EW, Inacio MC, Singh JA, Love R, Bini SA, Namba RS. (2015). Are There Modifiable Risk Factors for Hospital Readmission After Total Hip Arthroplasty in a US Healthcare System? *Clin Orthop Relat Res*, 473(11): 3446-3455.

Sheth D, Cafri G, Inacio MC, Paxton E, Namba R. (2015). Anterior and Anterolateral Approaches for THA Are Associated With Lower Dislocation Risk Without Higher Revision Risk. *Clin Orthop Relat Res*, 473(11): 3401-3408.

### 2014

Allepuz A, Havelin L, Barber T, Sedrakyan A, Graves S, Bordini B, Hoeffel D, Cafri G, Paxton E. (2014). Effect of Femoral Head Size on Metal-on-HXLPE Hip Arthroplasty Outcome in a Combined Analysis of Six National and Regional Registries. *J Bone Joint Surg Am*, 96(Supplement 1): 12-18.

# Publications

## Total Joint Replacement

*continued from page 38*

Banerjee S, Cafri G, Isaacs AJ, Graves S, Paxton E, Marinac-Dabic D, Sedrakyan A. (2014). A Distributed Health Data Network Analysis of Survival Outcomes. *J Bone Joint Surg Am*, 96(Supplement 1): 7-11.

Bini SA, Inacio MC, Cafri G. (2014). Two-Day Length of Stay is Not Inferior to 3 Days in Total Knee Arthroplasty with Regards to 30-Day Readmissions. *J Arthroplasty*, 30(5): 733–738.

Comfort T, et al. (2014). International comparative evaluation of fixed-bearing non-posterior-stabilized and posterior- stabilized total knee replacements. *J Bone Joint Surg Am*, 96(Supplement 1): 65-72.

Furnes O, Paxton E, Cafri G, Graves S, Bordini B.; Comfort T, Rivas MC, Banerjee S, Sedrakyan A. (2014). Distributed Analysis of Hip Implants Using Six National and Regional Registries: Comparing Metal-on-Metal with Metal-on-Highly Cross-Linked Polyethylene Bearings in Cementless Total Hip Arthroplasty in Young Patients. *J Bone Joint Surg Am*, 96(Supplement 1): 25- 33.

Graves S, et al. (2014). International comparative evaluation of knee replacement with fixed or mobile-bearing posterior- stabilized prostheses. *J Bone Joint Surg Am*, 96(Supplement 1): 59-64.

Inacio MC, Kritz-Silverstein D, Raman R, Macera CA, Nichols JF, Shaffer RA, Fithian DC. (2014). The Risk of Surgical Site Infection and Re-Admission in Obese Patients Undergoing Total Joint Replacement Who Lose Weight Before Surgery and Keep It Off Post-Operatively. *Bone Joint J*, 96-B(5): 629-35.

Inacio MCS, Paxton EW, Fisher D, Li R, Barber T, Singh J. (2014). Bariatric Surgery Prior to Total Joint Arthroplasty May Not Provide Dramatic Improvements in Post-Arthroplasty Surgical Outcomes. *J Arthroplasty*, 29(7): 1359–1364.

Inacio MCS, Silverstein DK, Raman R, Macera CA, Nichols JF, Shaffer RA, Fithian D. (2014). Weight Patterns Before and After Total Joint Arthroplasty and Characteristics Associated with Weight Change. *Perm J*, 18(1): 25-31.

Khatod M, Cafri G, Namba RS, Inacio M, Paxton EW. (2014). Risk Factors for Total Hip Arthroplasty Aseptic Revision. *J Arthroplasty*, 29(7): 1412–1417.

Miric A, Inacio M, Namba RS (2014). The Effect of Chronic Kidney Disease on Total Hip Arthroplasty. *J Arthroplasty*, 29(6): 1225-30.

Miric A, Inacio MC, Kelly MP, Namba RS. (2014). Can Total Knee Arthroplasty Be Safely Performed Among Nonagenarians? An Evaluation of Morbidity and Mortality Within a Total Joint Replacement Registry. *J Arthroplasty*, 29(8): 1635–1638.

Miric A, Inacio MC, Namba RS. (2014). Can Total Knee Arthroplasty Be Safely Performed in Patients With Chronic Renal Disease? *Acta Orthopaedica*, 85(1): 71-78.

Namba R, et al. (2014). International comparative evaluation of knee replacement with fixed or mobile non-posterior- stabilized implants. *J Bone Joint Surg Am*, 96(Supplement 1): 52-58.

Namba RS, Inacio MCS, Cafri G. (2014). Increased Risk of Revision For High Flexion Total Knee Replacement With Thicker Tibial Liners. *Bone Joint J*, 96(2): 217-23.

Paxton E, Cafri G, Havelin L, Stea S, Palliso F, Graves S, Hoeffel D, Sedrakyan A. (2014). Risk of Revision Following Total Hip Arthroplasty: Metal-on-Conventional Polyethylene Compared with Metal-on-Highly Cross-Linked Polyethylene Bearing Surfaces: International Results from Six Registries. *J Bone Joint Surg Am*, 96(Supplement 1): 19-24.

# Publications

## Total Joint Replacement

*continued from page 39*

Sedrakyan A, Graves S, Bordini B, Pons M, Havelin L, Mehle S, Paxton E, Barber T, Cafri G. (2014). Comparative Effectiveness of Ceramic-on-Ceramic Implants in Stemmed Hip Replacement: A Multinational Study of Six National and Regional Registries. *J Bone Joint Surg Am*, 96(Supplement 1): 34-41.

Sedrakyan A, Paxton E, Graves S, Love R, Marinac-Dabic D. (2014). National and international postmarket research and surveillance implementation. *J Bone Joint Surg Am*, 96(Supplement 1): 1-6.

Singh JA, Inacio MC, Namba RS, Paxton EW. (2014). Rheumatoid arthritis is associated with higher 90-day hospital readmission rates compared to osteoarthritis after hip or knee arthroplasty: A cohort study. *Arthritis Care & Research*, 67(5): 718-724.

Stea S, et al. (2014). Multinational comprehensive evaluation of the fixation method used in hip replacement: interaction with age in context. *J Bone Joint Surg Am*, 96(Supplement 1): 42-51.

## 2013

Adams AL, Paxton EW, Wang JQ, Johnson ES, Bayliss EA, Ferrara A, Nakasato C, Bini SA, Namba RS. (2013). Surgical Outcomes of Total Knee Replacement According to Diabetes Status and Glycemic Control, 2001 to 2009. *J Bone Joint Surg Am*, 95(6): 481-7.

Bini S, Khatod M, Cafri G, Chen Y, Paxton EW. (2013). Surgeon, Implant, and Patient Variables May Explain Variability in Early Revision Rates Reported for Unicompartmental Arthroplasty. *J Arthroplasty*, 95(24): 2195-202.

Bini S, Khatod M, Inacio M, Paxton EW. (2013). Same-Day Versus Staged Bilateral Total Knee Arthroplasty Poses No Increase in Complications in 6672 Primary Procedures. *J Arthroplasty*, 29(4): 694-7.

Bini SA, Chen Y, Khatod M, Paxton EW. (2013). Does Pre-Coating Total Knee Tibial Implants Affect the Risk of Aseptic Revision? *Bone Joint J*, 95-B(3): 367-70.

Inacio M, Kritz-Silverstein D, Raman R, Macera C, Nichols J, Shaffer R, Fithian D. (2013). The Impact of Pre-operative Weight Loss on Incidence of Surgical Site Infection and Readmission After Total Joint Arthroplasty. *J Arthroplasty*, 29(3): 458-64.

Inacio MC, Ake CF, Paxton EW, Khatod M, Wang C, Gross TP, Kaczmarek RG, Marinac-Dabic D, Sedrakyan A. (2013). Sex and Risk of Hip Implant Failure: Assessing Total Hip Arthroplasty Outcomes in the United States. *JAMA Internal Medicine*, 173(6): 435-41.

Inacio MC, Cafri G, Paxton EW, Kurtz SM, Namba RS. (2013). Alternative Bearings in Total Knee Arthroplasty: Risk of Early Revision Compared to Traditional Bearings. *Acta Orthopaedica*, 84(2): 145-52.

Mohan V, Inacio M, Namba R, Sheth D, Paxton E. (2013). Monoblock All-Polyethylene Tibial Components Have a Lower Risk of Revision than Metal-Backed Modular Components: An Analysis of 27657 Primary Total Knee Arthroplasties. *Acta Orthopaedica*, 84(6): 530-6.

Namba R, Inacio MC, Paxton EW. (2013). Risk Factors Associated with Deep Surgical Site Infections After Primary Total Knee Arthroplasty: An Analysis of 56,216 Knees. *J Bone Joint Surg Am*, 95(9): 775-82.

Namba RS, Cafri G, Khatod M, Inacio MC, Brox TW, Paxton EW. (2013). Risk Factors for Total Knee Arthroplasty Aseptic Revision. *J Arthroplasty*, 28(8 Suppl): 122-7.

## 2012

Inacio MC, Kritz-Silverstein D, Paxton EW, Fithian DC. (2012). Do Patients Lose Weight After Joint Arthroplasty Surgery? A Systematic Review. *Clin Orthop Relat Res*, 471(1): 291-8.

Khatod M, Inacio M, Bini S. (2012). Short-Term Outcomes of Unresurfaced Patellas in Total Knee Arthroplasty. *J Knee Surg*, 26(2): 105-8.



# Publications

## Total Joint Replacement

*continued from page 40*

Khatod M, Inacio MC, Bini SA, Paxton EW. (2012). Pulmonary Embolism Prophylaxis in More Than 30,000 Total Knee Arthroplasty Patients: Is There a Best Choice? *J Arthroplasty*, 27(2): 167-72.

Namba RS, Inacio MC, Paxton EW. (2012). Risk Factors Associated with Surgical Site Infection in 30,491 Primary Total Hip Replacements. *J Bone Joint Surg Br*, 94(10): 1330-8.

Namba RS, Inacio MCS, Paxton EW, Ake CF, Wang C, Gross TP, Marinac-Dabic D, Sedrakyan A. (2012). Risk of Revision for Fixed Versus Mobile-Bearing Primary Total Knee Replacements. *J Bone Joint Surg Am*, 94(21): 1929-35.

Paxton E, Ake C, Inacio M, Khatod M, Marinac-Dabic D, Sedrakyan A. (2012). Evaluation of Total Hip Arthroplasty Devices Using a Total Joint Replacement Registry. *Pharmacoepidemiol Drug Saf*, 21(S2): 53-9.

## 2011

Inacio MC, Paxton EW, Chen Y, Harris J, Eck E, Barnes S, et al. (2011). Leveraging Electronic Medical Records for Surveillance of Surgical Site Infection in a Total Joint Replacement Population. *Infect Control Hosp Epidemiol*, 32(4): 351-9.

Khatod M, Inacio M, Bini S, Paxton E. (2011). Prophylaxis Against Pulmonary Embolism in Patients Undergoing Total Hip Arthroplasty. *J Bone Joint Surg Am*, 93(19): 1767-72.

Namba R, Inacio M, Paxton E, Robertsson O, Grave S. (2011). The Role of Registry Data in the Evaluation of Mobile Bearing Total Knee Arthroplasty. *J Bone Joint Surg Am*, 93(Supplement 3): 48-50.

Paxton E, Furnes O, Inacio M, Fenstad A, Namba R, Havelin L. (2011). Comparison of the Norwegian Knee Arthroplasty Register and a United States Arthroplasty Registry. *J Bone Joint Surg Am*, 93(Supplement 3): 20-30.

Robertsson O, Mendenhall S, Paxton E, Inacio M, Graves S. (2011). Challenges in Prosthesis Classification. *J Bone Joint Surg Am*, 93(Supplement 3): 72-75.

Sedrakyan A, Paxton E, Marinac-Dabic D. (2011). Stages and Tools for Multinational Collaboration: The Perspective from the Coordinating Center of the International Consortium of Orthopaedic Registries. *J Bone Joint Surg Am*, 93(Supplement 3): 76-80.

Sedrakyan A, Paxton E, Phillips C, Namba R, Funahashi T, Barber T, Sculco T, Padgett D, Wright T, Marinac-Dabic D. (2011). The International Consortium of Orthopaedic Registries: Overview and Summary. *J Bone Joint Surg Am*, 93(Supplement 3): 1-12.

## 2010

Bini SA, Fithian DC, Paxton LW, Khatod MX, Inacio MC, Namba RS, et al. (2010). Does Discharge Disposition After Primary Total Joint Arthroplasty Affect Readmission Rates? *J Arthroplasty*, 25(1): 114-7.

Paxton EW, Inacio MC, Khatod M, Yue EJ, Namba RS. (2010). Kaiser Permanente National Total Joint Replacement Registry: Aligning Operations with Information Technology. *Clin Orthop Relat Res*, 468(10): 2646-63.

Paxton EW, Namba RS, Maletis GB, Khatod M, Yue EJ, Davies M, et al. (2010). A Prospective Study of 80,000 Total Joint and 5000 Anterior Cruciate Ligament Reconstruction Procedures in a Community-Based Registry in the United States. *J Bone Joint Surg Am*, 92(2): 117-32.

## 2009

Namba RS, Chen Y, Paxton EW, Slipchenko T, Fithian DC. (2009). Outcomes of Routine Use of Antibiotic-Loaded Cement in Primary Total Knee Arthroplasty. *J Arthroplasty*, 24(6): 44-7.

# Publications

## Total Joint Replacement

*continued from page 41*

### 2008

Khatod M, Inacio M, Paxton EW, Bini SA, Namba RS, Burchette RJ, et al. (2008). Knee Replacement: Epidemiology, Outcomes, and Trends in Southern California: 17,080 Replacements from 1995 through 2004. *Acta Orthopaedica*, 79(6): 812-9.

Paxton EW, Inacio M, Slipchenko T, Fithian DC. (2008). The Kaiser Permanente National Total Joint Replacement Registry. *Perm J*, 12(3): 12-6.

### 2007

Namba RS, Inacio M. (2007). Early and Late Manipulation Improve Flexion After Total Knee Arthroplasty. *J Arthroplasty*, 22(6): 58-61.

### 2006

Khatod M, Barber T, Paxton E, Namba R, Fithian D. (2006). An Analysis of the Risk of Hip Dislocation with a Contemporary Total Joint Registry. *Clin Orthop Relat Res*, 447.

### 2005

Namba RS, Paxton L, Fithian DC, Stone ML. (2005). Obesity and Perioperative Morbidity in Total Hip and Total Knee Arthroplasty Patients. *J Arthroplasty*, 20(7 Suppl 3): 46-50.

Paxton L, Fithian DC. (2005). Outcome Instruments for Patellofemoral Arthroplasty. *Clin Orthop Relat Res*(436): 66-70.

### 2004

Fithian DC, Paxton EW, Cohen AB. (2004). Indications in the Treatment of Patellar Instability. *J Knee Surg*, 17(1): 47-56.

Fithian DC, Paxton EW, Post WR, Panni AS. (2004). Lateral Retinacular Release: A Survey of the International Patellofemoral Study Group. *J Arthroscopy*, 20(5): 463-8.

Fithian DC, Paxton EW, Stone ML, Silva P, Davis DK, Elias DA, White LM. (2004). Epidemiology and Natural History of Acute Patellar Dislocation. *Am J Sports Med*, 32(5): 1114-21.

### 2003

Sachs RA, Smith JH, Kunej M, Paxton L. (2003). Does Anticoagulation Do More Harm Than Good?: A Comparison of Patients Treated Without Prophylaxis and Patients Treated with Low-Dose Warfarin After Total Knee Arthroplasty. *J Arthroplasty*, 18(4): 389-95.

## Multi-Registry

### 2013

Campion TR, Jr., Johnson SB, Paxton EW, Mushlin AI, Sedrakyan A. (2013). Implementing Unique Device Identification in Electronic Health Record Systems: Organizational, Workflow, and Technological Challenges. *Med Care*, 52(1): 26-31.

Paxton E, Kiley M, Love R, Barber T, Funahashi T, Inacio M. (2013). Kaiser Permanente Implant Registries Benefit Patient Safety, Quality Improvement, Cost-Effectiveness: Innovation in Patient Safety and Quality at the National Level. *The Joint Commission Journal on Quality and Patient Safety*, 39(6): 246-52.

### 2012

Paxton E, Inacio M, Kiley M. (2012). The Kaiser Permanente Implant Registries Effect on Patient Safety, Quality Improvement, Cost Effectiveness, and Research Opportunities. *Perm J*, 16(2): 33-40.

### 2011

Kiley ML, Paxton E, Maas K. (2011). Registries Revolutionize Care and Outcomes. *Biomed Instrum Technol*, 45(2): 126-9.

# Publications

## Other Related Research

### 2017

Cafri G, Fan J. Between-within effects in survival models with cross-classified clustering: Application to the evaluation of the effectiveness of medical devices. *Stat Methods Med Res*, 2018; 27(1): 312-9.

Franklin PD, Chenok KE, Lavalee D, Love R, Paxton L, Segal C, Holve E. Framework to guide the collection and use of patient-reported outcome measures in the learning healthcare system. *eGems*, 2017; 5(1): 1-17.

### 2016

Inacio MC, Paxton EW, Dillon MT. (2016). Understanding Orthopaedic Registry Studies: A Comparison with Clinical Studies *J Bone Joint Surg Am*, 98 (1): 1-7.

Lavallee DC, Chenok KE, Love RM, Patersen C, Holve E, Segal CD, Franklin PD. (2016). Incorporating Patient-Reported Outcomes Into Health Care To Engage Patients And Enhance Care *Health Aff*, 35 (4): 575-82.

### 2015

Inacio MC, Chen Y, Paxton E, Namba R, Kurtz S, Cafri G. (2015). Statistics in Brief: An Introduction to the Use of Propensity Scores. *Clin Orthop Relat Res* 1-5.

### 2014

Le C, Guppy KH, Axelrod YV, Hawk MW, Silverthorn J, Inacio MC, Akins PT. (2014). Lower Complication Rates For Cranioplasty With Peri-Operative Bundle. *Clin Neurol Neurosurg*, 120 41-4.

### 2013

Csintalan R, Latt L, Fornalski S, Raiszadeh K, Inacio M, Fithian D. (2013). Medial Patellofemoral Ligament (MPFL) Reconstruction for the Treatment of Patellofemoral Instability. *J Knee Surg*, 27(2): 139-46.

### 2012

Maletis GB, Inacio MC, Reynolds S, Funahashi TT. (2012). Incidence of Symptomatic Venous Thromboembolism After Elective Knee Arthroscopy. *J Bone Joint Surg Am*, 94(8): 714-20.

### 2010

Harness NG, Inacio MC, Pfeil FF, Paxton LW, et al. (2010). Rate of Infection After Carpal Tunnel Release Surgery and Effect of Antibiotic Prophylaxis. *J Hand Surg*, 35(2): 189-96.

### 2005

Panni AS, Tartarone M, Patricola A, Paxton EW, Fithian DC. (2005). Long-term results of lateral retinacular release. *J Arthroplasty*, 21(5): 526-31.